

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (JSTO)**



# ABSTRACTS

## SOCIETY FOR NEUROSCIENCE

29TH ANNUAL MEETING • MIAMI BEACH, FLA. • OCTOBER 23-28, 1999

# 25

Part 1

Univ. of Minn.  
Bio-Medical  
Library

02 18 00

PART 1 OF 2

ischemia as compared to the control groups, non-ischemic (100%), sham-operated (100%), untreated ischemic (<5%), vehicle-treated (<5%; normal saline) ischemic rats. Animals which received the LD followed by a continuous IV infusion of  $\text{MgSO}_4$  at either 15, 30, 60 or 120 mg/kg/hr showed 30%, 80%, 15% and <5% neuronal preservation, respectively. The 90 mg/kg LD and 30 mg/kg/hr IV infusion was then administered 4, 8, 12 or 24 hours after ischemia and resulted in 80%, 70%, 55% and 35% neuronal preservation, respectively.

These results highlight: (i) a previously undocumented  $\text{MgSO}_4$  dose response, with higher doses being ineffective; and (ii) the neuroprotective potential for pre- and post-ischemic IV administration of  $\text{MgSO}_4$ .

Supported by ANRI/SCGH and RACS.

2364

#### **ISCHEMIC SEVERITY AND NEUROPROTECTION BY CDP-CHOLINE.**

**J. F. Hatcher, R. J. Dempsey\* and A. M. Rao.** Dept. of Neurological Surgery, Univ. of Wisconsin, Madison, WI 53792.

CDP-choline (CDPC) is a rate-limiting intermediate in the biosynthesis of phosphatidylcholine (PC), an important component of the neuronal membrane. The ability of CDPC to alter phospholipid metabolism has been shown to be an important function in experimental ischemia. Exogenous treatment with CDPC stimulates PC synthesis and prevents release of free fatty acids (FFA) especially arachidonic acid (AA) after ischemia/reperfusion. AA is one of the risk factors directly acting on neurons and its accumulation is greatest in vulnerable brain regions in ischemic injury. Here we report the neuroprotective effect of CDPC in hippocampal  $\text{CA}_1$  region after transient ischemia of gerbils. **Results:** Ischemia-reperfusion resulted in AA and leukotriene  $\text{C}_4$  accumulation, blood-brain barrier dysfunction, and edema; CDPC significantly attenuated all of these parameters. 5- or 10-min global ischemia followed by 6-d reperfusion resulted in  $88 \pm 7\%$  neuronal loss in the  $\text{CA}_1$  subfield of the hippocampus. CDPC (500mg/kg i.p.) administered just after 5-min ischemia and thereafter daily for 5 d significantly attenuated neuronal death to  $12 \pm 4\%$  ( $p < 0.05$  compared to untreated ischemic group). When ischemic duration was increased to 10-min, CDPC also significantly decreased neuronal death to  $31 \pm 6\%$  ( $p < 0.05$  compared to untreated ischemic group). CDPC administered as one dose (immediately after ischemia) did not provide significant protection, whereas two doses (at 0 and 1-d) provided slight neuroprotection. **Conclusions:** Accumulated AA after transient ischemia is converted to oxygenated metabolites through cyclooxygenase/lipoxygenase pathways, which are involved in BBB dysfunction, edema and neuronal loss. CDPC may protect neurons by preventing the membrane PC breakdown into AA and subsequent free radical-generating metabolites.

Funded by University of Wisconsin, NIH and Department of Veterans Affairs.

THIS PAGE BLANK (USPTO)

~~BEST AVAILABLE COPY~~